ed for ready absorption and slow perspiration, so that it derives from the dews and rain, what the earth fails to supply it.

The Gaultheria procumbens is remarkable for the different periods of producing its flowers and fruit. It is found in blossom not only in the early part of spring, but in the last weeks of summer, and the fruit is found ripe at corresponding periods. Whether this appearance is the product of different shoots, or whether the same stems blossom twice in a year, I am unable to say. I have, however, met with beds of the Gaultheria in full flower in August and September, quite as frequently as in May. I have also seen the fruit in the market at various periods of the summer, fall, and spring.

The plant takes its vulgar names from the fruit, and is denominated in different parts of the United States, *Partridge berry, Chequer berry, Box berry,* &c. Its domestic use has also given it the name of *Mountain tea.*

The genus Gaultheria is beautifully singular and distinct in its character, derived from the form of its fruit. The *calyx is five cleft, calyculated, or bibracteate at base. Corolla ovate. Capsule five celled, invested with the baccated calyx.*

The species *procumbens* has a *prostrate stem with ascending branches. Leaves in a terminal tuft, obovate with a few ciliate serratures. Flowers axillary.*

Class *Decandria*, order *Monogynia*, Natural orders *Bicornes Linn. Ericæ Juss.*

The stem, or as it might be called root of this plant is horizontal, woody, often a quarter of an inch in thickness. The branches are ascending, but a few inches high, round and somewhat downy. Leaves scattered, near the extremities of the branches, evergreen, coriaceous, shining, oval or obovate, acute at both ends, revolute at the edge, and furnished with a few small serratures, each terminating in a bristle. Flowers axillary, drooping, on round downy stalks. Outer calyx of two concave, heart shaped leafets, which may with perhaps more propriety be called bractes. Inner calyx monophyllous, white, cleft into five roundish subacute segments. Corolla white, urceolate, five angled, contracted at the mouth, the border divided into five short, reflexed segments, Filaments white, hairy, bent in a semicircular manner to accommodate themselves to the cavity between the corolla and germ. Anthers oblong, orange coloured, ending in two double horns, bursting outwardly, for their whole length above the filaments,